

## [SAO/NASA ADS Astronomy Abstract Service](#)

· [Find Similar Abstracts](#) (with [default settings below](#))

· [On-line Data](#)

· [Associated Articles](#)

· [Also-Read Articles](#) ([Reads History](#))

·

· [Translate This Page](#)

<b>Title:</b>		VizieR Online Data Catalog: BVI photometry of Galactic Cepheids (Tammann+, 2003)
<b>Authors:</b>		<a href="#">Tammann, G. A.</a> ; <a href="#">Sandage, A.</a> ; <a href="#">Reindl, B.</a>
<b>Publication:</b>		VizieR On-line Data Catalog: J/A+A/404/423. Originally published in: 2003A&A...404..423T
<b>Publication Date:</b>		06/2003
<b>Origin:</b>		<a href="#">SIMBAD</a>
<b>Keywords:</b>		Stars: variable, Photometry: UBVRI
<b>Comment:</b>		table1.dat 324x84 Intrinsic colors of Galactic fundamental-mode; Cepheids
<b>Bibliographic Code:</b>		<a href="#">2003yCat..34040423T</a>

### Abstract

321 Galactic fundamental-mode Cepheids with good B,V,I photometry by Berdnikov et al. (2000, Cat. ) and with homogenized color excesses  $E(B-V)$  based on Fernie et al. (1995IBVS.4148....1F) are used to determine their period-colour relation in the range  $0.4 < B < 4.17$ ,  $R_V = 3.17$ , and  $R_I = 1.89$ . The two sets define two independent P-L relations which agree very well in slope. They are therefore combined into a single mean Galactic P-L relation. (1 data file).